DEPARTMENT of ENVIRONMENTAL SERVICES Water Supply & Pollution Control Division - Biology Bureau

LAKE TROPHIC DATA

MORPHOMETRIC:

| Lake: TODD LAKE | Lake Area (ha): | 68.07 |
|------------------------------------|------------------------------------|----------|
| Town: NEWBURY | Maximum depth (m): | 6.1 |
| County: Merrimack | Mean depth (m): | 2.2 |
| River Basin: Merrimack | Volume (m³): | 1466500 |
| Latitude: 43°16'45" N | Relative depth: | 0.7 |
| Longitude: 71°58'05" W | Shore configuration: | 1.74 |
| Elevation (ft): 670 | Areal water load (m/yr) | : 1.17 |
| Shore length (m): 5100 | Flushing rate (yr ⁻¹): | 0.50 |
| Watershed area (ha): 155.4 | P retention coeff.: | 0.88 |
| <pre>% watershed ponded: 6.1</pre> | Lake type: natura | ıl w/dam |

| BIOLOGICAL: | 30 December 1991 | 17 July 1991 | |
|------------------------------------|------------------|----------------------|--|
| DOM. PHYTOPLANKTON (% TOTAL) #1 | ASTERIONELLA 80% | CHRYSOSPHAERELLA 70% | |
| #2 | | MELOSIRA 10% | |
| #3 | | | |
| PHYTOPLANKTON ABUNDANCE (cells/mL) | | 535 | |
| CHLOROPHYLL-A (µg/L) | | 4.74 | |
| DOM. ZOOPLANKTON (% TOTAL) #1 | KELLICOTTIA 47% | NAUPLIUS LARVAE 30% | |
| #2 | CODONELLA 17% | POLYARTHRA 22% | |
| #3 | | KELLICOTTIA 21% | |
| ROTIFERS/LITER | 47 | 180 | |
| MICROCRUSTACEA/LITER | 5 | 154 | |
| ZOOPLANKTON ABUNDANCE (#/L) | 70 | 344 | |
| VASCULAR PLANT ABUNDANCE | | Common/Abun | |
| SECCHI DISK TRANSPARENCY (m) | | 3.0 | |
| BOTTOM DISSOLVED OXYGEN (mg/L) | 8.1 | 1.8 | |
| BACTERIA (fecal col., #/100 ml) #1 | | < 10 | |
| #2 | | < 10 | |
| #3 | | | |

SUMMER THERMAL STRATIFICATION:

stratified

Depth of thermocline (m): 4.8 Hypolimnion volume (m^3) : None Anoxic volume (m^3) : None

| CHEMICAL: | Lake: TODD LAKE Town: NEWBURY | | | | |
|--------------------------|----------------------------------|-------|--------|---|--------|
| | 30 December 1991 17 July 1991 | | | | |
| DEPTH (m) | 2.0 | 4.0 | 2.0 | | 5.0 |
| pH (units) | 6.4 | 6.2 | 7.1 | | 6.5 |
| A.N.C. (Alkalinity) | 5.2 | 5.3 | 6.7 | | 7.3 |
| NITRATE NITROGEN | 0.02 | 0.03 | < 0.05 | | < 0.05 |
| TOTAL KJELDAHL NITROGEN | | | | | |
| TOTAL PHOSPHORUS | 0.003 | 0.005 | 0.011 | - | 0.014 |
| CONDUCTIVITY (µmhos/cm) | 40.3 | 40.4 | 45.1 | | 46.0 |
| APPARENT COLOR (cpu) | 18 | 22 | 18 | | 21 |
| MAGNESIUM | | | 0.54 | | |
| CALCIUM | | | 2.8 | | |
| SODIUM | | | 4.5 | | |
| POTASSIUM | | | 0.58 | | |
| CHLORIDE | 3 | 3 | 5 | | 5 |
| SULFATE | 6 | 5 | 3 | | 3 |
| TN : TP | | | | | |
| CALCITE SATURATION INDEX | | , | 2.9 | | |


All results in mg/L unless indicated otherwise

TROPHIC CLASSIFICATION: 1991

| D.O. | S.D. | PLANT | CHL | TOTAL | CLASS |
|------|------|-------|-----|-------|-------|
| ** | 3 | 4 | 1 | 8 | Meso. |

COMMENTS:

- 1. Also known as Todd Pond, Todd's Pond, Lake Todd and Todd's Lake.
- 2. In 1991 the Water Resources Division determined that Todd Lake was a natural pond over 10 acres, raised by damming, and thus a great pond. The Lake Todd Association does not accept that determination; they contend the pond is an artificial impoundment.
- 3. No public access.
- 4. This pond was previously surveyed and classified in 1977. The pond was eutrophic in 1977, but that determination included the use of a 1938 Fish & Game dissolved oxygen value. There was little change in water quality between the two years, although the subjective evaluation of plant abundance suggests that plants were somewhat less abundant in 1991 (this conclusion does not hold if the two plant maps are compared).



FIELD DATA SHEET

LAKE: TODD LAKE

DATE: 07/17/91

TOWN: NEWBURY

WEATHER: SUNNY & BREEZY


| PTH M) | TEMP (°C) | *DISSOLVED OXYGEN | OXYGEN SATURATION |
|-----------|--------------|----------------------|----------------------|
| 0.1 | 24.0 | 8.8 | 103 % |
| 1.0 | 24.0 | 8.6 | 101 % |
| 2.0 | 23.9 | 8.4 | 97 % |
| 3.0 | 22.5 | 8.2 | 93 % |
| 4.0 | 20.5 | 6.4 | 70 % |
| 5.0 | 15.2 | 1.8 | 18 % |
| 6.0 | 11.5 | 1.8 | 16 % |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| W | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

SECCHI DISK (m): 3.0 COMMENTS:

BOTTOM DEPTH (m): 6.2

TIME: 1245

*Dissolved oxygen values are in mg/L



AQUATIC PLANT SURVEY

| LAK | E: TODD LAKE | TOWN: NEWBURY | DATE: 07/17/91 |
|-----|---------------------------|-------------------------|----------------|
| Key | PLAN | ADIDIDANGE | |
| Key | GENERIC | COMMON | ABUNDANCE |
| P | Pontederia cordata | Pickerelweed | Common |
| U | Utricularia | Bladderwort | Scattered |
| N | Nymphaea | White water lily | Common |
| Y | Nuphar | Yellow water lily | Common |
| s | Sparganium | Bur reed | Common |
| В | Brasenia schreberi | Water shield | Scattered |
| W | Potamogeton | Pondweed | Common |
| F | Nymphoides cordatum | Floating heart | Common |
| g | | Bottom growth | Abundant |
| f | Chlorophyceae | Filamentous green algae | Scattered |
| D | Decodon verticillatus | Swamp loosestrife | Sparse |
| 0 | Cephalanthus occidentalis | Buttonbush | Sparse |
| т | Typha | Cattail | Sparse |
| M | Myriophyllum humile | Water milfoil | Sparse |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |
| | | | |

OVERALL ABUNDANCE: Common/Abun

GENERAL OBSERVATIONS:

- 1. Plants were particularly dense at the northern end of the pond.
- 2. Milfoil fragments were observed but no rooted plants.
- 3. Rhizosolenia (30%) was the dominant genus of wholewater plankton. The dominant classes were greens (35%) and diatoms (35%).